Nonlinear Analysis: Modelling and Control, Vol. VV, No. n, 1–NN
https://doi.org/10.15388/namc.2026.VV.XXXXXX

Author's version
November 3, 2026

Title of your article*

First Author[a†] **, Second Author**[b‡] **, Third Author**[b‡⊠]

[a]Institution a: Institute of Data Science and Digital Technologies, Faculty of Mathematics and Informatics, Vilnius University, Lithuania[ROR]
first.author@somewhere.host

[b]Institute of Applied Mathematics, Vilnius University, Lithuania[ROR]
second.author@somewhere.host; third.author@somewhere.host

Received: / Revised: / Published online:

Abstract. Abstract Text. A short abstract describing the research done, methodology, and achieved results is to be presented. The abstract should contain approximately 100 words. The volume of the article is up to 20 pages in NA journal style. The journal recognizes review articles as special ones. These articles (if any) will occupy the starting positions in the journal and may contain more than 20 pages. Text in article have to follow a few simple guidelines: complex mathematical expressions have to be justified (like in the excerpt below), algorithms have to be presented in the style of `alltt`, *postscript specials* have to be absent in file format presenting images. The enumeration of references within any article has to be organized in alphabetic order (see the sample below). References have to be distinguished between journal articles, collective works, and books and presented in BibTeX `NAMCplain` style

Keywords: keyword1; keyword2; keyword3.
AMS Subject Classification: ams1; ams2.

1 Introduction or the first section

This[1] is a sample demonstration of the requirements for articles submitted to the journal "Nonlinear Analysis: Modelling and Control". For visuality, we quote random portions of published articles.

1.1 Examples of citation

Citation of [3, article], [1, book], [2, proceedings]. Bibliography entries are included at the beginning of the *.tex file. When editing these entries, it is necessary to delete the 'x.bib' file before compiling the LaTeX document (this step is not required when working in Overleaf).

*This research was supported by grant No. xxxx.
†The Author suported by
‡The Author suported by
[1]Introduction text

© 2026 The Author(s).

2 Examples of equations, algorithms, figures

Let us consider one-way coupled chaotic systems of the following general form (master-slave configurations or systems with a skew product structure):

$$\dot{X} = F(X), \qquad \dot{Y} = G(Y, X). \tag{2.1}$$

Here $X \equiv \{x_1, x_2, \dots, x_d\}$ is a d-dimensional state vector of the driving system, and $Y \equiv \{y_1, y_2, \dots, y_r\}$ is an r-dimensional state vector of the response system. F and G define the vector fields of the driving and response systems.

2.1 Complex mathematical expressions

$$\begin{aligned} 1 \leqslant \bigl|P(1)\bigr| &= |a_d| \prod_{j=k+1}^{d} \alpha_j \prod_{j=k+1}^{d} \bigl(\alpha_j^{-1} - 1\bigr) \prod_{j=1}^{d} (\alpha_j - 1) \\ &\leqslant \frac{|a_d a_0|}{M(P)} \left(\left(\frac{M(P)}{|a_0|} \right)^{1/(d-k)} - 1 \right)^{d-k} \left(\left(\frac{M(P)}{|a_d|} \right)^{1/k} - 1 \right)^{k} \\ &= M(P)^{-1} \bigl(M(P)^{1/(d-k)} - |a_0|^{1/(d-k)} \bigr)^{d-k} \bigl(M(P)^{1/k} - |a_d|^{1/k} \bigr)^{k} \\ &\leqslant M(P)^{-1} \bigl(M(P)^{1/(d-k)} - 1 \bigr)^{d-k} \bigl(M(P)^{1/k} - 1 \bigr)^{k}, \end{aligned} \tag{2.2}$$

$$\begin{aligned} \Omega &= \bigl\{(r,z)\colon\ 0 < r < R,\ 0 < z < H\bigr\}, \\ \Omega_0 &= \bigl\{(r,z)\colon\ r^2 + z^2 < R_0^2,\ z > 0\bigr\}, \\ \Gamma_1 &= \bigl\{(r,0)\colon\ 0 \leqslant r \leqslant R\bigr\}, \\ \Gamma &= \bigl\{(0,z)\colon\ 0 < z \leqslant H\bigr\} \cup \bigl\{(R,z)\colon\ 0 < z \leqslant H\bigr\} \\ &\quad \cup \bigl\{(r,H)\colon\ 0 < r < R\bigr\} \cup \Gamma_1, \end{aligned} \tag{2.3}$$

$$u_{ij} = u(r_i, z_j, t), \qquad v_{ij} = v(r_i, z_j, t), \qquad d_{\alpha,ij} = d_\alpha(r_i, z_j),$$

$i = 0, 1, \dots, N$, $j = 0, 1, \dots, M_\alpha$, $\alpha = 1, 2$.

GS guarantees that the asymptotic dynamics of the response system is independent of its initial conditions and is completely determined by the driving system. Geometrically, this implies a collapse of the overall evolution onto a stable synchronization manifold $M = \{(X, Y)\colon\ \Phi(X) = Y\}$ in the full phase space of the two systems $X \oplus Y$. It is easy to show that the linear stability of the identity manifold $Y' = Y$ in the extended phase space $X \oplus Y \oplus Y'$ is equivalent to the linear stability of the manifold $M = \{(X, Y)\colon\ \Phi(X) = Y\}$ in the original $X \oplus Y$ phase space.

2.2 The following excerpt demonstrates presentation of an algorithm

```
Input: A list of segmentation parameters and the
       corresponding C1, C2 and C3 values.

Output: A selection of the optimum segmentation and its
       parameters.

 BEGIN
```

```
        Search the input list for local minima in C2,
        and list the local minima into L;

        Search L and locate the minimum of C1 + 10 C3^2,
        present this element of L as the result;
 END
```

2.3 Just another excerpt presenting image

In the works of S. Fučík, this spectrum was studied first as an object related to "slightly" nonlinear problems. The interest in these type of problems grew also in connection with the theory of suspension bridges (see, e.g., [6]). From the mathematical point of view, the Fučík equation became a source of numerous investigations generalizing and refining the results by Fučík.

The Fučík equation with Sturm–Liouville conditions has similar structure of spectrum, and the description of it can be found in [4] or [5].

Completely different spectrum was obtained for the problem composed of equation

$$x'' = -\mu x^+ + \lambda x^- \tag{2.4}$$

with nonlocal conditions

$$x(0) = 0, \qquad \int_0^1 x(s)\,\mathrm{d}s = 0. \tag{2.5}$$

A recent paper [8] deals with the problem composed of equation (2.4) with conditions

$$x'(0) = 0, \qquad \int_0^1 x(s)\,\mathrm{d}s = 0. \tag{2.6}$$

Some branches of the spectra for these two problems are depicted in Fig. 1.




Figure 1. The spectrum for problem (2.4), (2.5) and for problem (2.4), (2.6).

3 Conclusion

Acknowledgment. We would like to thank dr. N. Sergejeva for the excerpts of article [7].

Author contributions. All authors (F.A., S.A. and T.A.) have contributed as follows: Methodology, S.A.; formal analysis, F.A., S.A and T.A.; software, F.A. and T.A.; validation, F.A. and T.A.; writing–original draft preparation, F.A.; writing–review and editing, S.A.. and T.A.; visualization, F.A. and T.A. All authors have read and agreed to the published version of the manuscript.

Conflicts of interest. The authors declare no conflicts of interest.

References

1. A. Author, *Title of Book*, Volume 3 of *B*, Publisher, Address, 5 edition, 2027, https://doi.org/10.15388/namc.2024.30.42210.
2. F. Author, S. Author, Title of proceedings, in Editor (Ed.), *BOOKTITLE*, Volume 3 of *A*, publisher, address, month 2027, pp. 1–6, https://doi.org/10.15388/namc.2022.30.42210. note
3. F. Author, S. Authors, T. Author, Title of Article, *Journal*, **1**(3):101–120, 2027, https://doi.org/10.15388/namc.2023.30.42210.
4. E.N. Dancer, On the Dirichlet problem for weakly non-linear elliptic partial differential equations, *Proc. R. Soc. Edinb., Sect. A, Math.*, **76**:283–300, 1977, https://doi.org/10.1017/S0308210500019648.
5. S. Fučík, *Solvability of Nonlinear Equations and Boundary Value Problems*, Reidel, Dordrecht, 1980, https://doi.org/10.1080/1023629033000062934.
6. A.C. Lazer, P.J. McKenna, Large-amplitude periodic oscillations in suspension bridges: Some new connections with nonlinear analysis, *SIAM Rev.*, **32**(4):537–578, 1990, http://www.jstor.org/stable/2030894.
7. N. Sergejeva, The Fučík spectrum for nonlocal BVP with Sturm–Liouville boundary condition, *Nonlinear Anal. Model. Control*, **19**(3):503–513, 2014, https://doi.org/10.15388/NA.2014.3.13.
8. N. Sergejeva, On the Fučík type problem with the integral nonlocal boundary conditions, *Electron. J. Qual. Theory Differ. Equ.*, **21**(1):1–11, 2016, https://doi.org/10.14232/ejqtde.2016.8.21.